arXiv:1208.4636v1 [math.NT] 22 Aug 2012

Three-Dimensional Solvable Artin Representations Ramified at One Prime

Jonah Leshin

August 24, 2012

Abstract

We classify the possibilities for the fixed field of the kernel of an irreducible three-dimensional Artin representation of $\mathbb{Q}$ with solvable image ramified at one prime by using the classification of the finite irreducible subgroups of $\mathrm{PGL}_3(\mathbb{C})$. This allows us to bound the number of such representations with given Artin conductor.

1 Introduction

The Langlands program gives a correspondence between Galois representations and automorphic representations. The analytic objects corresponding to two dimensional Artin representation are modular forms of weight one and Maass forms of weight zero, which correspond to odd and even two dimensional Artin representations, respectively. Specifically, Deligne and Serre [2] showed that given a cuspidal modular form of weight 1, level N, and Nebentypus character ϵ with normalized $(a_1 = 1)$ Fourier expansion $\sum a_n q^n$, there exists an Artin representation

$$\rho : \mathrm{G}_{\mathbb{Q}} := \mathrm{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \to \mathrm{GL}_2(\mathbb{C})$$

of conductor N such that (via ρ) the trace of the conjugacy class of the Frobenius at p Frob_p in $\mathrm{G}_{\mathbb{Q}}$ is a_p and the determinant of Frob_p is $\epsilon(p)$. Conversely, by the Langlands-Tunnell theorem [1], given a two dimensional Artin representation ρ, if we insist that ρ has solvable image, then there is known to exist an automorphic form f such that f gives rise to ρ in the manner described above. Using this correspondence, Wong [14], building on work of Duke [4], bounded the number of two dimensional Artin representations of $\mathbb{Q}$ with Artin conductor N in terms of N.

Less is known in the case of three-dimensional Artin representations. The goal of this paper is to bound the number of three-dimensional irreducible Artin representations with solvable image and given conductor. It is known that such representations come from automorphic representations of $\mathrm{GL}_3(\mathbb{A}_{\mathbb{Q}})$, where $\mathbb{A}_{\mathbb{Q}}$ denotes the adèles of $\mathbb{Q}$. We use purely algebraic techniques for our bounds, which thus give bounds for the number of automorphic representations of $\mathrm{GL}_3(\mathbb{A}_{\mathbb{Q}})$ with particular properties.

The main results of this paper, which are made more precise in Theorems 1, 2, 4, and 5 are:

- The number of imprimitive three-dimensional Artin representations of $\mathrm{G}_{\mathbb{Q}}$ with conductor dividing p^m is at most Cmp^{m+1}, where C is an explicitly given constant.
- The number of primitive three-dimensional Artin representations of $\mathrm{G}_{\mathbb{Q}}$ with conductor dividing p^m is at most $Dp^{\frac{m}{3}+50}$, where D is an explicitly given constant.

The paper is divided into two main parts. The first part looks at imprimitive representations. We use the fact that any imprimitive three-dimensional representation is monomial to bound the number of all imprimitive representations in one fell swoop. The second part deals with primitive representations, where we do a case by case analysis of the three groups that can occur as the projective image of a primitive three-dimensional Artin representation.

2 Definitions and Preliminaries

Let V be a finite dimensional vector space over an arbitrary field, and let G be a group. A representation $\rho : G \to \mathrm{GL}(V)$ is called *imprimitive* if there exist proper subspaces $V_1, \dots, V_m$ of V such that $V = \oplus_{i=1}^m V_i$ and $\rho(G)$ permutes the V_i. If such a decomposition of V does not exist, ρ is called *primitive.* If ρ is imprimitive and the corresponding V_i are of dimension 1, then ρ is said to be *monomial* (this is equivalent to the standard definition of monomial, which is that there exists a subgroup $H \leq G$ and a linear character χ of H such that ρ is induced from χ). Thus monomial representations of dimension greater than 1 are imprimitive, and the converse is true if the dimension of the representation is prime. A group is said to be *monomial* if all its irreducible representations are monomial.

An Artin representation of a number field F is a continuous homomorphism

$$\rho : \mathrm{Gal}(\bar{F}/F) \to \mathrm{GL}_n(\mathbb{C}).$$

We take the topology on $\mathrm{GL}_n(\mathbb{C})$ to be discrete, so Artin representations necessarily have finite image. The conductor $N(\rho)$ of ρ is defined to be the product of its local conductors, whose definition we now recall. Let $\mathfrak{p}$ be a prime of F. For the $\mathfrak{p}$-part $N_{\mathfrak{p}}(\rho)$ of $N(\rho)$, consider the restriction $\rho_{\mathfrak{p}}$ of ρ to $\mathrm{G}(\bar{F}_{\mathfrak{p}}/F_{\mathfrak{p}})$ (if E/F is an extension of fields, we will often write $\mathrm{G}(E/F)$ in place of $\mathrm{Gal}(E/F)$ to ease notation). The representation ρ_p factors through a finite quotient $\mathrm{G}(K/F_{\mathfrak{p}})$ of $\mathrm{G}(\bar{F}_{\mathfrak{p}}/F_{\mathfrak{p}})$. Let g_i denote the order of the ith ramification group G_i of $\mathrm{G}(K/F_{\mathfrak{p}})$. We then define $N_{\mathfrak{p}}(\rho)$ to be $\mathfrak{p}$ raised to the power

$$\sum_{i \geq 0} \frac{g_i}{g_0} \mathrm{dim} V/V^{G_i},$$

where V is the representation space for ρ and V^{G_i} is the subspace of elements fixed by G_i. For the remainder of this paper we make the following hypothesis on all Artin representations:

$$F = \mathbb{Q} \quad \text{and} \quad n = 3.$$

The natural projection $\pi : \mathrm{GL}_3(\mathbb{C}) \to \mathrm{PGL}_3(\mathbb{C})$ will always be denoted by π. A cyclic group of order n will be denoted C_n, and $Z(G)$ denotes the center of a group G. If L/K is a Galois extension of fields, we say E is a central extension of L/K if E/K is Galois and $\mathrm{G}(E/L) \leq Z\big(\mathrm{G}(E/K)\big)$.

The symbol E_ρ will denote the fixed field of the kernel of one of our three-dimensional Artin representations ρ, and F_ρ will denote the fixed field of the kernel of $\pi \circ \rho$ (when the context is clear, the subscript ρ may be dropped). Thus ρ descends to an injection $\mathrm{G}(E_\rho/\mathbb{Q}) \hookrightarrow \mathrm{GL}_3(\mathbb{C})$, which by abuse of notation we also denote by ρ, and $Z\big(\mathrm{G}(E/\mathbb{Q})\big) = \mathrm{G}(E/F)$, a cyclic group.

We say an Artin representation ρ is *ramified at p* if the extension $E_\rho/\mathbb{Q}$ is ramified at p. In sections 3 and 4, all representations are assumed to be unramified outside $p\infty$, where p is a finite prime and ∞ is the real place of $\mathbb{Q}$. To avoid special cases (most of which are trivial), we will assume $p \neq 2, 3$.

In our computations, we will wish to bound the number of C_a or $C_a \times C_a$ extensions of a number field K with given ramification conditions. We do this by first finding the a-rank of the maximal extension L of K with the given ramification conditions. Let $r_{a,b}(x)$ be the number of $(C_a)^b$ extensions of K contained in a Galois extension L, where $\mathrm{G}(L/K)$ has a-rank x. We will consider the cases when $a = 2$ or 3, although the following formulas hold whenever a is prime:

$$r_{a,1} = \frac{a^x - 1}{a - 1}, \quad r_{a,2} = \frac{(a^x - 1)(a^x - a)}{(a^2 - 1)(a^2 - a)}.$$

When we have an extension L/K of number fields, with $L/K, L/\mathbb{Q}$, and $K/\mathbb{Q}$ all Galois and $L/\mathbb{Q}$ ramified only at p, we will write $e(L/K)$ to denote the ramification from K to L of the primes above p in K. We will say an extension of fields is unramified if and only if it is unramified at all finite primes. The ring of integers of a number field K will be denoted by $\mathfrak{O}_K$ and its absolute discriminant by d_K. We let h_K denote the class number of K, $\mathrm{Cl}(K)$ the ideal class group of K, $\mathrm{Cl}^{\mathfrak{m}}(K)$ the ray class group of K modulo $\mathfrak{m}$, H_K the Hilbert class field of K, and $K^{\mathfrak{m}}$ the ray class field of K of modulus $\mathfrak{m}$. If $\mathfrak{m}$ is a product of finite primes, we will write $\mathfrak{m}\infty$ to denote the modulus that is the product of $\mathfrak{m}$ multiplied by all the real places of K.

3 Imprimitive Representations

In this section we assume that $\rho : \mathrm{G}_{\mathbb{Q}} \to \mathrm{GL}_3(\mathbb{C})$ is imprimitive. Since ρ is monomial, there exists a field L of degree 3 over $\mathbb{Q}$ and a character $\chi : \mathrm{G}_L \to \mathbb{C}^*$ such that $\rho = \chi_L^{\mathbb{Q}}$, where $\chi_L^{\mathbb{Q}}$ denotes the induced representation of χ to $\mathrm{G}_{\mathbb{Q}}$.

Suppose first that $L \subset \mathbb{Q}(\zeta_p)$ and let $\mathfrak{p}$ be the prime of L above p. It follows from the Corollary to Proposition 4 of [13] that

$$v_p(N(\rho)) = v_{\mathfrak{p}}(N(\chi)) + 2.$$

Each non-trivial $\chi \in \mathrm{Hom}(\mathrm{G}_L, \mathbb{C}^*)$ can be induced to an irreducible ρ. Since G_L is normal in $\mathrm{G}_{\mathbb{Q}}$, it follows from Mackey Decomposition [9] that if $\rho = \chi_L^{\mathbb{Q}}$, then there are three distinct

χ's in $\mathrm{Hom}(\mathrm{G}_L, \mathbb{C}^*)$ that give ρ when induced up to $\mathrm{G}_\mathbb{Q}$. Let

$$c_m = \#\{\rho : \rho = \chi_L^\mathbb{Q}, N(\rho) \mid p^m\} \text{ and } d_m = \#\{\chi : N(\chi) \mid \mathfrak{p}^m\}, \text{ so } c_m = \frac{d_{m-2} - 1}{3}.$$

We will first bound d_m. By class field theory, any χ with $N(\chi) \mid \mathfrak{p}^m$ factors through $\mathrm{G}(L^{\mathfrak{p}^m}/L)$, where $L^{\mathfrak{p}^m}$ is the ray class field of L modulo $\mathfrak{p}^m$. Class field theory gives the exact sequence

$$\mathfrak{O}_L^* \to (\mathfrak{O}_L/\mathfrak{p}^m\infty)^* \to \mathrm{Cl}^{\mathfrak{p}^m\infty}(L) \to \mathrm{Cl}(L) \to 1, \tag{1}$$

where $(\mathfrak{O}_L/\mathfrak{p}^m\infty)^*$ is defined to be $(\mathfrak{O}_L/\mathfrak{p}^m)^* \times \langle \pm 1\rangle^3$, the exponent 3 coming from the 3 real places of L. The image of $\mathfrak{O}_L^*$ in $(\mathfrak{O}_L/\mathfrak{p}^m\infty)^*$ has order at least two, so we obtain

$$d_m = |\,\mathrm{G}(L^{\mathfrak{p}^m\infty}/L)| \le 4h_L p^{m-1}(p-1) < 4\left(\frac{1}{2}p\right)p^{m-1}(p-1) = 2p^m(p-1),$$

where the first inequality follows from (1), and the second inequality from [10]. Thus we obtain

Theorem 1. *Notation as above,*

$$c_m < \frac{2p^{m-1}}{3}.$$

For the remainder of this section, suppose that $L \not\subseteq \mathbb{Q}(\zeta_p)$, or equivalently, that $L/\mathbb{Q}$ is not Galois. By the following lemma, p must split in L as $\mathfrak{p}_1^2\mathfrak{p}_2$.

Lemma 1. *Let $L/\mathbb{Q}$ be a degree 3 extension ramified at a single finite prime $p \neq 2, 3$. Then $L/\mathbb{Q}$ is Galois if and only if p is totally ramified in L.*

Proof. The only if part is clear. For the converse, suppose $L/\mathbb{Q}$ is totally ramified at p and let $\mathfrak{p}$ be the prime of L above p. Let $L_\mathfrak{p}$ be the localization of L at $\mathfrak{p}$ and $\mathfrak{O} = \mathfrak{O}_{L_\mathfrak{p}}$ the ring of integers in $L_\mathfrak{p}$. Let $f(x)$ be the minimal polynomial for $\alpha \in \mathfrak{O}$, where α generates $\mathfrak{O}$ as a polynomial ring over $\mathbb{Z}_p$. We may assume $\alpha \in \mathfrak{O}_L$. The p-part of the discriminant $\mathrm{disc}(f)$ of f equals the local discriminant $d_{L_\mathfrak{p}/\mathbb{Q}_p}$, which has p-part p^2 ($p \neq 3$), the same as the p-part of the global discriminant $d_{L/\mathbb{Q}}$. So we can write $\mathrm{disc}(f) = \epsilon p^2 D$, with $\epsilon \in \{\pm 1\}$ and $p \nmid D$. The Galois closure K of $L/\mathbb{Q}$ is ramified only at p and is given by $K = L\big(\sqrt{\mathrm{disc}\, f}\big) = L\big(\sqrt{\epsilon D}\big)$. For $p \neq 2$, this is only possible if $\sqrt{\epsilon D} \in L$, i.e. $L/\mathbb{Q}$ is Galois. □

By the Corollary to Proposition 4 of [13], we have

$$N(\chi_L^\mathbb{Q}) = N_{L/\mathbb{Q}}(N(\chi))d_L.$$

It follows that if $N(\chi) = \mathfrak{p}_1^{a_1}\mathfrak{p}_2^{a_2}$, then $N(\rho) = p^{a_1+a_2}p$. Let $c_m = c_{m,L}$ be the number of imprimitive ρ with conductor dividing p^m that are induced from G_L. To estimate c_m, we need to estimate

$$\#\{\chi : N(\chi) \mid \mathfrak{p}_1^{a_1}\mathfrak{p}_2^{a_2}, a_1 + a_2 \le m - 1\}.$$

Thus we need to bound $\sum_{a_1+a_2=m-1} |\,\mathrm{G}(L^{\mathfrak{p}_1^{a_1}\mathfrak{p}_2^{a_2}\infty}/L)|$. We have

$$\sum_{a_1+a_2=m-1} |\,\mathrm{G}(L^{\mathfrak{p}_1^{a_1}\mathfrak{p}_2^{a_2}\infty}/L)| \leq 4h_L \sum_{a_1+a_2=m-1} (p^{a_1}-1)(p^{a_2}-1) < 4h_L m p^{m-1},$$

where the first inequality follows from (1). Since $L/\mathbb{Q}$ is not Galois, G_L is not normal in $\mathrm{G}_{\mathbb{Q}}$, so it is possible that there may be only one $\chi \in \mathrm{Hom}(\mathrm{G}_L, \mathbb{C}^*)$ that gives ρ when induced up to $\mathrm{G}_{\mathbb{Q}}$. So we have

$$c_m < 4h_L m p^{m-1} - 1. \tag{2}$$

It remains to bound h_L and the number n_p of non-Galois fields L of degree 3 over $\mathbb{Q}$, ramified only at $p\infty$. We do the latter first. Let L be such a number field. The Galois closure K of $L/\mathbb{Q}$ is an S_3 extension of $\mathbb{Q}$ ramified only at $p\infty$. By Lemma 1, p splits as $\mathfrak{p}_1^2\mathfrak{p}_2$ in L. Let $\mathfrak{q}$ be a prime of K above $\mathfrak{p}_2$. We have

$$e(\mathfrak{q}/p) = e(\mathfrak{q}/\mathfrak{p}_2)e(\mathfrak{p}_2/p) = e(\mathfrak{q}/\mathfrak{p}_2) \leq 2. \tag{3}$$

For any primes $\mathfrak{q}_i, \mathfrak{q}_j$ of K above p, $e(\mathfrak{q}_i/p) = e(\mathfrak{q}_j/p)$, and the quantity is at least 2, so by (3), it equals 2. Thus $K/\mathbb{Q}(\sqrt{p^*})$ is unramified, where $\mathbb{Q}(\sqrt{p^*})$ is the subfield of K of degree 2 over $\mathbb{Q}$. Therefore, L and its two Galois conjugates contained in K combine to give one unramified C_3 extension of $\mathbb{Q}(\sqrt{p^*})$. By [11], we have $h_{\mathbb{Q}(\sqrt{p^*})} < \frac{22.2p}{\pi^2}$. Ramification at the infinite primes of $\mathbb{Q}(\sqrt{p^*})$ can only contribute a power of 2 to the degree of the maximal abelian unramified (at all finite primes) extension of $\mathbb{Q}(\sqrt{p^*})$. Since we are interested in the 3-rank of this extension, we may ignore the potential ramification at the infinite primes. We have the obvious bound that the 3-rank of $\mathrm{Cl}\left(\mathbb{Q}(\sqrt{p^*})\right)$ is less than

$$\log_3\left(\frac{22.2p}{\pi^2}\right),$$

from which it follows that

$$n_p < 3r_{3,1}\left(\log_3\left(\frac{22.2p}{\pi^2}\right)\right),$$

which is less than

$$\frac{3\cdot 11.1p}{\pi^2}.$$

Putting this last bound together with (2), and using the bound $h_L < \frac{22.2p}{\pi^3}$ (which follows from [11]), we obtain

Theorem 2. *The number of imprimitive ρ with $\rho = \chi_L^{\mathbb{Q}}$ and $L/\mathbb{Q}$ non-Galois is less than*

$$\left(\frac{2957.04mp^{m+1}}{\pi^5}\right).$$

Remark 1. *It is possible to obtain a slightly better bound by using a slightly better bound for the* 3*-rank of* $\mathrm{Cl}\left(\mathbb{Q}(\sqrt{p^*})\right)$*. See, for example, [12].*

4 Primitive Representations

For this section, suppose that $\rho : \mathrm{G}_{\mathbb{Q}} \to \mathrm{GL}_3(\mathbb{C})$ is primitive. We continue with our assumption that ρ is unramified outside $p\infty$. Since ρ is primitive, the image of $\pi \circ \rho$ in $\mathrm{PGL}_3(\mathbb{C})$ is isomorphic to one of the following groups [3]:

$$P_1 := \big(C_3 \times C_3\big) \rtimes C_4, \quad P_2 := \big(C_3 \times C_3\big) \rtimes Q_8, \quad P_3 := \big(C_3 \times C_3\big) \rtimes \mathrm{SL}_2(\mathbb{F}_3),$$

where Q_8 is the quaternion group of order 8. We can view P_1 as a subgroup of P_2, and P_2 as a subgroup of P_3.

We claim that $\mathrm{Im}(\pi \circ \rho)$ can not be P_2. If $\mathrm{G}(F_\rho/\mathbb{Q}) = P_2$, then F_ρ (and thus E_ρ) has a subfield L with $L/\mathbb{Q}$ Galois and $\mathrm{G}(L/\mathbb{Q}) \cong C_2 \times C_2$. But there is no such field ramified only at one prime ($p \neq 2$) of $\mathbb{Q}$.

In the next two subsections, we consider the cases $\mathrm{G}(F_\rho/\mathbb{Q}) = P_1$ and $\mathrm{G}(F_\rho/\mathbb{Q}) = P_3$. For each case, we first fix a number field F with $\mathrm{G}(F/\mathbb{Q}) = P_i$ and bound the number of representations ρ with $F_\rho = F$. We then bound the number of possible such F– i.e. F such that $\mathrm{G}(F/\mathbb{Q}) = P_i$ and $F/\mathbb{Q}$ is unramified outside $p\infty$.

4.1 Representations with Projective Image Isomorphic to P_1

We count the number of Artin representations ρ unramified outside $p\infty$ with projective image equal to P_1, i.e. $\mathrm{G}(F_\rho/\mathbb{Q}) \cong P_1$. Let L denote the fixed field of $C_3 \times C_3 \leq P_1$. Since L is unramified outside $p\infty$, it is the unique degree four subfield of $\mathbb{Q}(\zeta_p)$. Writing $[E : F] = n$, we have the following tower of fields.

$$\begin{tikzcd}
E \arrow[d, dash, "C_n"] \\
F \arrow[d, dash, "C_3 \times C_3"] \\
L \arrow[d, dash, "C_4"] \\
\mathbb{Q}
\end{tikzcd}$$

Lemma 2. *Let K be a subfield of $\mathbb{Q}(\zeta_p)$, and let $\mathfrak{p}$ be the unique prime of K above the rational prime p. Then $K^{\mathfrak{p}\infty} = H_K(\zeta_p)$.*

Proof. Certainly $H_K(\zeta_p) \subseteq K^{\mathfrak{p}\infty}$. Suppose the containment is strict. Then $K^{\mathfrak{p}\infty}/H_K(\zeta_p)$ is totally ramified at the primes of $H_K(\zeta_p)$ above p. It follows that $K^{\mathfrak{p}\infty}/\mathbb{Q}(\zeta_p)$ is tamely ramified at the prime $(1 - \zeta_p)$ of $\mathbb{Q}(\zeta_p)$ above p. But such an extension can not exist since $\mathbb{Q}(\zeta_p)^{(1-\zeta_p)} = H_{\mathbb{Q}(\zeta_p)}$ [8]. $\square$

We have $F \subseteq L^{\mathfrak{p}}$, where $\mathfrak{p}$ is the unique prime of L above p. Suppose that the ramification degree $e = e(F/L)$ of $\mathfrak{p}$ in F is 3 or 9. Then by Lemma 2, the ramification in F/L must be obtained by adjoining the unique degree 3 ($e = 3$) or 9 ($e = 9$) subfield of $\mathbb{Q}(\zeta_p)$ to L. It follows that $|P_1'| = [F : F \cap \mathbb{Q}^{ab}] = 3$ or 1 if $e = 3$ or 9, respectively. This is a contradiction since $P_1' = C_3 \times C_3$. Therefore, F/L is unramified and $F \subseteq H_L$.

We have the following lemma of Tate [5]:

Lemma 3. (Tate [5]) *For each prime p, let I_p be the inertia group of p in $\mathrm{G}_{\mathbb{Q}_p}$, which we view as a fixed subgroup of $\mathrm{G}_{\mathbb{Q}}$. Let $\tilde{\rho} : \mathrm{G}_{\mathbb{Q}} \to \mathrm{PGL}_n(\mathbb{C})$ be a projective representation. Suppose that for each p, there exists a lift $\rho_p : \mathrm{G}_{\mathbb{Q}_p} \to \mathrm{GL}_n(\mathbb{C})$ of $\tilde{\rho}|_{\mathrm{G}_{\mathbb{Q}_p}}$ with $\rho_p|_{I_p}$ trivial for all but finitely many p. Then there exists a unique lift $\rho : \mathrm{G}_{\mathbb{Q}} \to \mathrm{GL}_n(\mathbb{C})$ of $\tilde{\rho}$ such that $\rho|_{I_p} = \rho_p|_{I_p}$ for all p.*

Returning to our notation before Lemma 3, since $F_\rho/\mathbb{Q}$ is unramified outside p, for all primes $l \neq p$, $\tilde{\rho}|_{\mathrm{G}_{\mathbb{Q}_l}}$ factors through a cyclic extension of $\mathbb{Q}_l$. We may thus lift $\tilde{\rho}|_{\mathrm{G}_{\mathbb{Q}_l}}$ to a map $\rho_l : \mathrm{G}_{\mathbb{Q}_l} \to \mathrm{GL}_3(\mathbb{C})$ such that the fixed field of the kernel of $\tilde{\rho}|_{\mathrm{G}_{\mathbb{Q}_l}}$ is the same as the fixed field of the kernel of ρ_l; in particular, both fixed fields are unramified at l.

Since the ideal $\mathfrak{p}$ of L is principal and F_ρ/L is unramified, $\mathfrak{p}$ splits completely in F_ρ. Thus the kernel of the fixed field of $\tilde{\rho}|_{\mathrm{G}_{\mathbb{Q}_p}}$ is a totally tamely ramified extension of $\mathbb{Q}_p$, so cyclic. We may therefore lift $\tilde{\rho}|_{\mathrm{G}_{\mathbb{Q}_p}}$ to a map ρ_p such that the kernel of the fixed field of ρ_p is the same as the kernel of the fixed field of $\tilde{\rho}|_{\mathrm{G}_{\mathbb{Q}_p}}$.

Putting this local data together, Lemma 3 now tells us that there exists a lift ρ of $\tilde{\rho}$ such that the ramification in $F_\rho/\mathbb{Q}$ is the same as the ramification in $E_\rho/\mathbb{Q}$ (i.e. E_ρ is unramified at all primes $l \neq p$ and the ramification degree at p in $E_\rho/\mathbb{Q}$ is the same as the ramification degree at p in $F_\rho/\mathbb{Q}$).

We now come to the main result of this section.

Proposition 1. *Let $\rho : \mathrm{G}_{\mathbb{Q}} \to \mathrm{GL}_3(\mathbb{C})$ be an Artin representation unramified outside $p\infty$ with projective image isomorphic to P_1. Let F be the fixed field of $\ker \pi \circ \rho$. Let $\mathbb{Q}(\zeta_{p_3})$ denote the maximal subfield of $\mathbb{Q}(\zeta_p)$ of 3-power degree over L, and let $\mathbb{Q}(\zeta_{p_{3'}})$ be the composite of all subfields of $\mathbb{Q}(\zeta_{p^\infty})$ with degree prime-to-3 over L. Then the fixed field of $\ker \rho$ either*

i) *lies between E and $E\mathbb{Q}(\zeta_{p_{3'}})$, where E is the unique non-trivial unramified central extension of $F/\mathbb{Q}$, and $[E : F] = 3$, or*

ii) *lies between E_3' and $E_3'\mathbb{Q}(\zeta_{p_{3'}})$, where $E_3' \nsubseteq F\mathbb{Q}(\zeta_{p_3})$ is one of two possible degree 3 ramified extensions of FM that is central over $F/\mathbb{Q}$, where M is a subfield of $\mathbb{Q}(\zeta_{p_3})$.*

Proof. There are several things to prove. First we show that if $F/\mathbb{Q}$ has a non-trivial unramified central extension, then that extension is unique. We will use the following theorem.

Theorem 3. (Fröhlich [6]) *Let L/K be Galois and let N be a central extension of L/K. Then there is a surjective map*

$$H_2(\mathrm{G}(L/K), \mathbb{Z}) \twoheadrightarrow \mathrm{G}\left(N/L(N \cap K^{ab})\right),$$

and there exists a central extension N' of L/K such that

$$H_2(\mathrm{G}(L/K), \mathbb{Z}) \xrightarrow{\sim} \mathrm{G}\left(N'/L(N' \cap K^{ab})\right).$$

Moreover, if $K = \mathbb{Q}$, one can take N' to be ramified at the same rational primes that L is ramified at.

In our situation, $H_2(\mathrm{G}(F/\mathbb{Q}), \mathbb{Z}) = C_3$. Suppose that E_1 and E_2 are distinct non-trivial unramified central extensions of $F/\mathbb{Q}$. We have $F(E_i \cap \mathbb{Q}^{ab}) \subseteq E_i \cap F\mathbb{Q}^{ab}$, so by Theorem 3, $C_3 \twoheadrightarrow \mathrm{G}\left(E_i/E_i \cap F\mathbb{Q}^{ab}\right), i = 1, 2$. Since E_i/F and F/L are unramified, $E_i \cap F\mathbb{Q}^{ab} = F$. Suppose $E_1 \neq E_2$. Then E_1E_2/F is an unramified central extension of $F/\mathbb{Q}$ with $E_1E_2 \cap F\mathbb{Q}^{ab} = F$ and $[E_1E_2 : F] = 9$, contradicting Theorem 3.

We now show the existence of a non-trivial unramified (over F) central extension of $F/\mathbb{Q}$. Let ρ' be a lift of $\pi \circ \rho$ such that the image of $\rho|_{I_p}$ has order 3, the existence of which is guaranteed by Lemma 3. Let E be the fixed field of $\ker \rho'$. Since F/L is unramified, the ramification in $E/\mathbb{Q}$ takes place in $L/\mathbb{Q}$, from which it follows that E/F is unramified and $E \cap F\mathbb{Q}^{ab} = F$. By Theorem 3, we must have $[E : E \cap F\mathbb{Q}^{ab}] = [E : F] \leq 3$. We can not have $E = F$ because the group P_1 does not have any irreducible three-dimensional representations, so $[E : F] = 3$. Thus E gives the desired extension.

Now let ρ be any representation for which $F_\rho = F$ and let E' be a candidate for the fixed field of $\ker \rho$. By Theorem 3, there exists $M' \subset \mathbb{Q}(\zeta_{p^\infty})$ such that $[E' : FM'] \leq 3$, and $E' \cap \mathbb{Q}^{ab} = M'$. We claim that $E' \neq FM'$. If E' were contained in FM', then we would have $\mathrm{G}(E/\mathbb{Q}) \cong P_1 \times C$, where C is cyclic. But since P_1 has no irreducible primitive three-dimensional representations, neither does $P_1 \times C$. So $[E' : FM'] = 3$.

Suppose first that E'/FM' is unramified. Being a composite of two central extensions of F, EM' is a central extension of F as well (see Figure 1).



Figure 1: Field diagrams for proof of Proposition 1.

If $E' \neq EM'$, then EE' is an unramified central extension of FM'. Since F/L is unramified, EE'/FM' being unramified implies that $EE' \cap FM'\mathbb{Q}^{ab} = FM'$, and $[EE' : FM'] = 9$, which contradicts Theorem 3. Suppose that $3 \mid [M' : \mathbb{Q}]$. Then EM'/F, having two intermediate subfields of degree 3 over F, is not cyclic. Thus we must have $M' \subseteq \mathbb{Q}(\zeta_{p_{3'}})$.

Suppose for the remainder of the proof that E'/FM' is ramified. Let E''/FM' be another degree 3 ramified extension. The extension $E'E''/FM'$ can not be totally ramified because if it were, the corresponding extension of local fields at primes above p would be tamely ramified but not cyclic. Therefore, FM' has a degree 3 unramified extension contained in $E'E''$. The field EM' is unramified over FM' and is central over $F/\mathbb{Q}$. If there were to exist another such field, the composite D of this field with EM' would be unramified over FM' and central over $F/\mathbb{Q}$, but we would have $[D : D \cap F\mathbb{Q}^{ab}] = 9$, contradicting Theorem 3. It follows that EM' is the unique degree 3 unramified extension of FM' contained in $E'E''$. To avoid contradicting Theorem 3, we must have $[E'E'' : E'E'' \cap F\mathbb{Q}^{ab}] = 3$. We have $\mathrm{G}(E'E''/FM') \cong C_3 \times C_3$, so there are 4 intermediate C_3 extensions of FM': E', E'', EM', and $E'E'' \cap F\mathbb{Q}^{ab}$. Of these, E'' is the only field besides E' that can be ramified over FM' and the fixed field of ρ.

Let $M_3' = M' \cap \mathbb{Q}(\zeta_{p_3})$. There exists a degree 3 extension E_3' of FM_3' such that $E' = E_3'M'$, and E_3'/FM_3' is ramified. This completes the proof of Proposition 1. ☐

We note the following, which is the existence of E from case i) of Proposition 1.

Corollary 1. *Notation as above, suppose that L admits an unramified $C_3 \times C_3$ extension F. Then $F/\mathbb{Q}$ admits a degree three unramified central extension.*

The next goal is to count the number of representations $\mathrm{G}_{\mathbb{Q}} \to \mathrm{GL}_3(\mathbb{C})$ with projective image isomorphic to P_1 and given prime power conductor. We must have $p \equiv 1 \pmod 4$, and we will also assume (as is necessary for such a representation to exist), that the 3-rank k of the class group $\mathrm{Cl}(L)$ of L is at least 2. Then L admits $r_{3,2}(k)$ unramified extensions F with $\mathrm{G}(F/L) = C_3 \times C_3$.

Let $G = \mathrm{G}(E/\mathbb{Q})$, where E is as in case i) of Proposition 1. Since the 3-Sylow subgroup $\mathrm{G}(E/L) := A$ of G is normal in G, by Hall's Theorem on solvable groups [7], G splits as $A \rtimes C_4$. If A is abelian, then G is metabelian (i.e. the second commutator group of G is trivial), which gives a contradiction since metabelian groups are monomial [9], and ρ is primitive. Suppose that A is isomorphic to the non-trivial semi-direct product $C_9 \rtimes C_3$. A generator of C_4 acts on $G/Z(G)$ as an automorphism of order 4 and thus must act on A as an automorphism of order a multiple of 4. But $|\operatorname{Aut}(C_9 \rtimes C_3)| = 54$, so no such automorphism exists. So A must be the other non-abelian group of order 27, $(C_3 \times C_3) \rtimes C_3$. The group G has 8 irreducible three-dimensional representations. Recall the definition of the Artin conductor $N(\rho)$ of a representation ρ from Section 1. We have $G_0 = C_4$, $G_i = 0, i \geq 1$. One finds that for 2 of the 8 aforementioned representations, $\dim V^{G_0} = 0$, and for the other 6, $\dim V^{G_0} = 1$, where $V \cong \mathbb{C}^3$ is the representation space. So we get two representations with conductor p^3 and 6 with conductor p^2.

Suppose first that we are in case i) of Proposition 1, so the fixed field of the kernel of a representation ρ is of the form EM, with $M \subset \mathbb{Q}(\zeta_{p_{3'}})$. Let $b = [EM : E]$. The irreducible three-dimensional representations of $\mathrm{G}(EM/\mathbb{Q})$ are given by $\rho \otimes \chi$, where χ is a character of $\mathrm{G}(M/L)$, of which there are b (note that $\mathrm{G}(EM/\mathbb{Q}) = \big((C_{3b} \times C_3) \rtimes C_3\big) \rtimes C_4)$. Thus the number of irreducible representations of ρ of dimension three with $\ker\rho$ having fixed field EM is $8b$.

Continuing with case i) of Proposition 1, we now fix a $C_3 \times C_3$ extension F of L and count the number of ρ for which $N(\rho) = p^m$ and $F_\rho = F$. Write $[M : \mathbb{Q}] = 4cp^n$, with $c \mid \frac{p-1}{4}$, so $b = p^n c$. Let G_i, with order g_i, denote the lower ramification groups of $\mathrm{G}(EM/\mathbb{Q})$. As soon as $M \not\subseteq E$, we get non-trivial central elements in G_i whenever $G_i \neq 1$. Such elements act by scalars and thus do not fix any non-zero subspace of V. So for such ρ, $v_p(N(\rho)) = \sum_{i\geq 0} \frac{g_i}{g_0} \cdot 3$. We must compute the g_i. We know $g_0 = 4cp^n, g_1 = p^n$. The group G_0 can be identified with $\mathrm{G}(M/\mathbb{Q})$, where $M \subseteq \mathbb{Q}(\zeta_{p^{n+1}})$. Write G'_i and G'^i for the lower and upper, respectively, ramification groups of $G' := \mathrm{G}\left(\mathbb{Q}(\zeta_{p^{n+1}})/\mathbb{Q}\right)$. As shown in [13, Chapter IV.4], for any $0 \leq k \leq n$,

$$|G'_i| = p^k \quad \text{for} \quad p^{n-k} \leq i \leq p^{n-k+1} - 1, \quad i \in \mathbb{Z}.$$

We also have

$$|G'^i| = p^{n+1-\lceil i \rceil}, i \in \mathbb{R}_{>0},$$

where $\lceil i \rceil$ denotes the least integer greater than or equal to i. Let $H = \mathrm{G}\left(\mathbb{Q}(\zeta_{p^n+1})/M\right)$ so that $G'/H \cong G_0$. If G^i is a p-group, which it is for all $i > 0$, then

$$G^i = (G'/H)^i = G'^i H/H = G'^i / G'^i \cap H = G'^i, \tag{4}$$

where the second equality is a property of upper ramification groups. The lower ramification groups are determined by the upper ramification groups, so (4) implies that $G_i = G'_i$ for all $i \geq 1$ (see [13, Chapter IV] for details about the upper and lower ramification groups). So $|G_i| = p^k$ for $p^{n-k} \leq i \leq p^{n-k+1} - 1$ and $0 \leq k \leq n$, and $|G_i| = 1$ for $i \geq p^n$. Thus if $\ker\rho$ has fixed field EM,

$$v_p\big(N(\rho)\big) = 3\left(1 + \frac{p-1}{4c} + \frac{p(p-1)}{4cp} + \cdots + \frac{p^{n-1}(p-1)}{4cp^{n-1}}\right) = 3 + 3n\frac{p-1}{4c},$$

and there are $8p^n c$ such representations.

We are now in a position to estimate the number $a_{m,i}$ of ρ for which we are in case i) of Proposition 1 and for which $N(\rho) = p^m$ and $F_\rho = F$. If $m \equiv 1$ or $2 \pmod 3$, then $a_{m,i} = 0$. Suppose $m \equiv 0 \pmod 3$. Let $d = \gcd(\frac{p-1}{4}, k)$, where $m = 3k + 3$. Then

$$a_{m,i} = 8p^k \frac{p-1}{4} + \cdots + 8p^{\frac{k}{d}} \frac{p-1}{4d} = 8 \sum_{j|d} p^{\frac{k}{j}} \frac{p-1}{4j}.$$

The first term in the sum, for example, corresponds to the case $c = \frac{p-1}{4}, n = k$, in which case there exist $8p^k \frac{p-1}{4}$ representations ρ with $v_p\big(N(\rho)\big) = 3 + 3n\frac{p-1}{4c} = 3 + 3k$.

For the total number $A_{m,F,i}$ of representations ρ in case i) of Proposition 1 with $F_\rho = F$ and $N(\rho) \mid p^m$, we have

$$A_{m,F,i} \sim \sum_{0 \leq k \leq \frac{m-3}{3}} 2p^{k+1} \sim 2p^{\frac{m}{3}},$$

where we are viewing $A_{m,F,i}$ as a function of p.

Suppose now that we are in case ii) of Proposition 1. We can write the fixed field of $\ker \rho$ as $E'M'$, with $M' \subset \mathbb{Q}(\zeta_{p_3'})$ and E' a degree 3 ramified extension of FM, with $M \subseteq \mathbb{Q}(\zeta_{p_3})$. Let $[M : L] = 3^a$. A similar analysis to that of case i) shows that $\mathrm{G}(E'/\mathbb{Q}) \cong B \rtimes C_4$, where $B \cong (C_{3^{a+1}} \times C_3) \rtimes C_3$, with C_4 acting trivially on $C_{3^{a+1}}$. The group $B \rtimes C_4$ has $8 \cdot 3^a$ irreducible three-dimensional representations and $\mathrm{G}(E'M'/\mathbb{Q})$ has $8p^nc$, where $[E'M' : \mathbb{Q}] = 108p^nc$, $(c,p) = 1$. Let G_i, with order g_i, be the lower ramification groups of $\mathrm{G}(E'M'/\mathbb{Q})$. For all $i \geq 0$, G_i has non-trivial central elements, so $\dim V/V^{G_i} = 3$ whenever $g_i \neq 1$. So again we obtain

$$v_p\left(N(\rho)\right) = \sum \frac{g_i}{g_0} \cdot 3, \tag{5}$$

and we have to compute the g_i. We have $g_0 = 12cp^n$. In case i), we were able to compute the ramification groups because we could identify G_0 with a cyclotomic extension. In this case, we know g_0. For $i \geq 1$, g_i is a p-group, and the following lemma allows us to compute that for all i, the p-part of g_i for case ii) is the same as that for case i).

Lemma 4. *Let $M \subseteq \mathbb{Q}_p(\zeta_p)$ ($p \neq 2$), and let F/M be a Galois extension of degree prime to p. Let $\alpha \in \mathbb{Q}_p(\zeta_{p^\infty})$ be such that $[M(\alpha) : M]$ is a power of p. Then for $i \geq 1$, the lower ramification groups of $M(\alpha)/M$ have the same order as (and are isomorphic to) the lower ramification groups of $F(\alpha)/F$.*

Proof. We have $\mathrm{G}(F(\alpha)/F) \cong \mathrm{G}(M(\alpha)/M)$ and by looking at ramification degrees, we must have $F(\alpha)/F$ totally ramified. That the g_i for $F(\alpha)/F$ equal the g_i for $M(\alpha)/M$ now follows from the details of the computation of the latter that can be found in [13, Chapter IV.4]. □

We are thus able to compute, analogously to case i),

$$v_p\big(N(\rho)\big) = 3 + 3n\frac{p-1}{12c}.$$

In the same way that we bounded $a_{m,i}$, we can now bound the number $a_{m,ii}$ of ρ for which we are in case ii) of Proposition 1 and for which $N(\rho) = p^m$ and $F_\rho = F$. By (5), $a_{m,ii} = 0$ unless $a_{m,ii} \equiv 0 \pmod 3$, so write $a_{m,ii} = 3k + 3$. A similar series of computations to those in case i) gives

$$a_{m,ii} = 8p^k\frac{p-1}{4} + \cdots + 8p^{\frac{k}{d}}\frac{p-1}{12d} = 8\sum_{j|d} p^{\frac{k}{j}}\frac{p-1}{12j}\ , \quad d = \gcd\left(\frac{p-1}{4}, k\right).$$

For a fixed F and M, we've seen that for case ii) there are two possible degree 3 extensions of FM, so we obtain

$$A_{m,F,ii} \sim \sum_{0 \leq k \leq \frac{m-3}{3}} \frac{4}{3}p^{k+1} \sim \frac{4}{3}p^{\frac{m}{3}}.$$

To bound the total number B_m of primitive representations ρ with $N(\rho) \mid p^m$ and $\mathrm{Im}(\pi \circ \rho) = P_1$, we need to multiply $A_{m,F,i} + A_{m,F,ii}$ by the number of unramified $C_3 \times C_3$

extensions of L, which, as previously discussed, is $r_{3,2}(k)$, where k is the 3-rank of $\mathrm{Cl}(L)$. From [11], the 3-part of h_L is less than $\frac{22.2p^3}{\pi^4}$, so $k \leq \log_3(\frac{22.2p^3}{\pi^4}) := R$.

We introduce the following (non-standard) notation. If A and B are both positive, real-valued functions of n, we write $A \lessapprox B$ to mean $\limsup_{n\to\infty} \frac{A(n)}{B(n)} \leq 1$. Using this notation, we obtain:

Theorem 4. *Notation as above,*

$$B_m \lessapprox (A_{m,F,i} + A_{m,F,ii}) r_{3,2}(R) \sim \frac{222}{3^5\pi^4} p^{\frac{m+9}{3}}.$$

4.2 Representations with Projective Image Isomorphic to P_3

Recall that $P_3 := \big(C_3 \times C_3\big) \rtimes \mathrm{SL}_2(\mathbb{F}_3)$. Using the notation from the previous section, we look to count the number of ρ with $\mathrm{G}(F_\rho/\mathbb{Q}) \cong P_3$ and given conductor. Our main tool will be the following proposition.

Proposition 2. *Let E_ρ be the fixed field of the kernel of a primitive three-dimensional Artin representation ρ with $N(\rho)$ a pth power and $\mathrm{Im}\,\pi \circ \rho = P_3$. If $M \subseteq \mathbb{Q}(\zeta_{p^\infty})$, then there are at most three degree 3 central extensions of $F_\rho M/\mathbb{Q}$ ramified over $F_\rho M$ and at most one that is unramified over $F_\rho M$. There exists $M \subseteq \mathbb{Q}(\zeta_{p^\infty})$ such that of the four possible degree 3 central extensions of $F_\rho M/\mathbb{Q}$, E_ρ is either an unramified extension or one of two of the three possible ramified extensions.*



Figure 2: Field diagram for Section 4.2.

Proof. We have $H_2(P_3, \mathbb{Z}) \cong C_3$. Let $F = F_\rho$ and let M be a number field contained in $\mathbb{Q}(\zeta_{p^\infty})$. We claim that $FM/\mathbb{Q}$ may admit a maximum of three distinct non-trivial central extensions E_i with $E_i \cap \mathbb{Q}^{ab} = M$. By Theorem 3, for each i, we have $[E_i : FM] = 3$. The compositum $\prod E_i$ is a central extension of $FM/\mathbb{Q}$. By Theorem 3, there exists a cyclotomic field M' containing M such that $|\,\mathrm{G}(\prod E_i/FM')| \leq 3$. This means $\prod E_i/FM$ has rank at most two, i.e. its Galois group is C_3 or $C_3 \times C_3$. Such a field extension has at most (in the $C_3 \times C_3$ case) four strictly intermediate extensions– three $E_i's$ and FM', and we must have $[M' : M] = 3$.

The group $\mathrm{G}(FM'/\mathbb{Q})$ has an abelian normal subgroup that is not cyclic, and such a group has no faithful primitive representations [7], so FM' can not be the fixed field of $\ker \rho$ (the same argument shows FN can not be the fixed field of $\ker \rho$ for any $N \subset \mathbb{Q}(\zeta_{p^\infty})$). Since we can not have a totally ramified $C_3 \times C_3$ extension of FM that is Galois over $\mathbb{Q}$, we must have either at least one of the $E_i's$ be unramified or have FM'/FM unramified. If FM'/FM were unramified, ramification considerations force $e := e(F/F_0)$ to be 3 or 9, where F_0 is the fixed field of the $C_3 \times C_3$ normal subgroup of P_3. This is impossible, however, since we would have $e(F/\mathbb{Q}) = 3e$, so the extension of $\mathbb{Q}_p$ given by localizing F at a prime above p would have a C_{3e} subgroup, and P_3 does not have such a subgroup. Thus one of the E_i's is unramified over FM. If two of the E_i's were unramified, the composite of these fields would give a $C_3 \times C_3$ unramified central extension of FM, forcing FM'/FM to be unramified by Theorem 3, which we saw gives a contradiction.

Thus, $FM/\mathbb{Q}$ may have up to four central extensions of degree 3, three of which (all but FM') are possibilities for E_ρ. This completes the proof since E_ρ is a degree 3 central extension of $FM/\mathbb{Q}$ for some $M \subset \mathbb{Q}(\zeta_{p^\infty})$. □

Remark 2. *Unlike in the P_1 case, we may not use Lemma 3 here to guarantee the existence of an unramified degree 3 central extension of F because the extension of $\mathbb{Q}_p$ given by localizing at a prime above p in F is not necessary cyclic, so we are not assured of a lift $\rho_p : \mathrm{G}_{\mathbb{Q}_p} \to \mathrm{GL}_3(\mathbb{C})$ of $\tilde{\rho}|_{\mathrm{G}_{\mathbb{Q}_p}}$ such that the fixed field of the kernel of ρ_p is unramified over the fixed field of the kernel of $\tilde{\rho}|_{\mathrm{G}_{\mathbb{Q}_p}}$.*

Let $L = F^{ab}$ be the degree 3 extension of $\mathbb{Q}$ contained in $\mathbb{Q}(\zeta_p)$. The field L then admits a $C_2 \times C_2$ extension, which is unramified by Lemma 2. We have seen that F/F_0 must be unramified, so $e(F/\mathbb{Q}) = 3$ or 6.

Continuing in the manner of the P_1 case, we now bound, for a fixed $F/\mathbb{Q}$ with $\mathrm{G}(F/\mathbb{Q}) \cong P_3$, the number of ρ for which $N(\rho) = p^m$ and $F_\rho = F$. Let $x = e(F/\mathbb{Q})/3$. Let $E = E_\rho$, so E is a degree three extension of FM for some cyclotomic field M. Given M, by Proposition 2 there are at most three possible choices for E. Write $[M : L] = p^n c$, with $c \mid \frac{p-1}{3}$. Let $\delta = 3$ or 1, depending on whether E/FM is ramified or unramified, respectively. So $e(E/\mathbb{Q}) = 3p^n cx\delta$. Computing similarly to the P_1 case, we find

$$v_p(N_\rho) = 3 + 3n\frac{p-1}{3cx\delta}.$$

There are 3 non-split central extensions of P_3 of order $3|P_3| = 648$, and each has 7 irreducible three-dimensional representations. Thus $\mathrm{G}(E/\mathbb{Q})$ has $7p^n c$ irreducible three-dimensional representations. We estimate the numbers $a_{m,F,nr}$ and $a_{m,F,ram}$ of representations ρ with $N(\rho) = p^m$, $F_\rho = F$, and E_ρ/F unramified, and ramified, respectively. We must have $3 \mid m$ so write $m = 3k + 3$. Proceeding as in the P_1 case, and assuming that for any M, FM may admit 3 distinct central extensions as candidates for E_ρ, one of which is unramified over

FM, we see that the leading terms in the expressions for $a_{m,F,nr}$ and $a_{m,F,ram}$ are

$$\frac{7p^k(p-1)}{3x} \text{ and } 2\cdot\frac{7p^k(p-1)}{9x}, \text{ respectively.} \tag{6}$$

As we did with P_1, we must now bound the number of fields F with $\mathrm{G}(F/\mathbb{Q}) \cong P_3$ and $F/\mathbb{Q}$ ramified only at $p\infty$. As above, let $L = F^{ab}$, the degree 3 subfield of $\mathbb{Q}(\zeta_p)$. The field L then admits a Q_8 extension F_0, which has 3 intermediate fields L_i', each of which is a C_2 extension of L, and unramified over L by Lemma 2. Let K be the composite of the L_i. The field L is totally real, so by the following lemma, K is as well.

Lemma 5. *Let k be a totally real field and let k/l be a Galois extension with group Q_8. Then the unique intermediate field that is degree 4 over l is totally real.*

Proof. Let $k' = k \cap \mathbb{R}$. If $k' = k$, we are done. Otherwise, $[k : k'] = 2$, and k' is the unique intermediate field of degree 4 over l. ☐

Returning to the notation before Lemma 5, K/L is unramified and K is totally real, so the number of choices for K is at most $r_{2,2}(k)$, where k is the 2-rank of of $\mathrm{Cl}(L)$. Using [10], we have $|\mathrm{Cl}(L)| < \frac{1}{2}p$, so $k < \log_2 p - 1$. The field F_0 is a degree 2 extension of K, and the extension F_0/K is at most tamely ramified, so $F_0 \subseteq K^{\mathfrak{p}\infty}$, where $\mathfrak{p}$ is the prime of L above p, viewed as an ideal of K. By class field theory $\mathfrak{p}$ is a product of 4 primes in K since it is principal in L. Using the exact sequence (1) from Section 3 applied to $\mathrm{Cl}^{\mathfrak{p}\infty}(K)$, we find that $\mathrm{G}(K^{\mathfrak{p}\infty}/K)$ has 2-rank at most $4 + 12 +$ the 2-rank of $\mathrm{Cl}(K)$. The order of $\mathrm{Cl}(K)$ is at most $\frac{22.2p^8}{\pi^{12}}$ [11]. Set $\beta = \log_2(2^{12}\cdot\frac{22.2p^8}{\pi^{12}})$. If F_0/K is unramified above p, there are at most

$$r_{2,1}(\beta)r_{2,2}(\log_2 p - 1) \tag{7}$$

choices for F_0 and if F_0/K is ramified above p, there are at most

$$\left(r_{2,1}(4+\beta) - r_{2,1}(\beta)\right) r_{2,2}(\log_2 p - 1). \tag{8}$$

The final step to bound the number of possible F_ρ is to look at unramified $C_3 \times C_3$ extensions of F_0 (which give our field F_ρ). This amounts to finding $r_{3,2}(l)$, where l is the 3-rank of $\mathrm{Cl}(F_0)$. Using [11] and the fact that ramification in the infinite places does not contribute to the 3-rank, we obtain

$$l \le \log_3 \frac{22.2p^{16}}{\pi^{24}} \tag{9}$$

if F_0/K is unramified, and

$$l \le \log_3 \frac{22.2p^{20}}{\pi^{24}} \tag{10}$$

if F_0/K is ramified.

We have $x = 1$ or 2, depending on whether F_0/K is ramified. To obtain our bound for the number a_m (where $m = 3k+3$) of ρ ramified only at $p\infty$ with $\mathrm{G}(F_\rho/\mathbb{Q}) \cong P_3$ and

$N(\rho) = p^m$, we need to combine (6) with (7) and (9), and combine (6) with (8) and (10). Doing so, we find that the leading term of a_m in p, which comes only from the $x = 2$ case, is bounded by

$$(8.6 \times 10^{-23})p^{k+51}.$$

Finally, to conclude the section on P_3, we find that for the number B_m of three-dimensional Artin representations with projective image P_3 and conductor dividing p^m, we have

Theorem 5. *Notation as above,*

$$B_m \lessapprox (8.6 \times 10^{-23})p^{\frac{m}{3}+50}.$$

5 Comparison to the Two-Dimensional Case

For comparison to the two-dimensional case, we look at [14]. Looking at Theorem 3 in [14] and multiplying by $\phi(p^m) = p^{m-1}(p-1)$ to account for all possible χ, we see that the bounds for the number of two dimensional representations with conductor p^m and projective image isomorphic to A_4 or S_4 (and conjecturally for A_5) is on the order of $a_m = Cp^{2m-\epsilon} \log^4 p^m$ for $\epsilon = \frac{1}{6}, \frac{1}{8}$, or $\frac{1}{12}$ and C a constant, both ϵ and C depending on the projective image of the representation. Taking $p \to \infty$ for fixed m, we find that the bounds in [14] are better than our bounds for the imprimitive case. By going through the proofs of Theorems 4 and 5 in this paper, we may bound the number of ρ with $N(\rho) \mid p^m$, $\operatorname{Im} \pi \circ \rho = P_1$, and with $N(\rho)|p^m$, $\operatorname{Im} \pi \circ \rho = P_3$, respectively, as $m \to \infty$ for fixed p. These bounds can be given by easily computable constants multiplied by the bounds given in Theorems 4 and 5. Therefore, our bounds as $m \to \infty$ are stronger than those in [14], and our bounds are stronger as either p or m tends to ∞ for the primitive case. Of course, Wong's bounds hold for any conductor N, not only for prime power conductors.

References

[1] Gary Cornell, Joseph H. Silverman, and Glenn Stevens, editors. *Modular forms and Fermat's last theorem.* Springer-Verlag, New York, 1997. Papers from the Instructional Conference on Number Theory and Arithmetic Geometry held at Boston University, Boston, MA, August 9–18, 1995.

[2] Pierre Deligne and Jean-Pierre Serre. Formes modulaires de poids 1. *Ann. Sci. École Norm. Sup. (4)*, 7:507–530 (1975), 1974.

[3] Igor V. Dolgachev. Finite subgroups of the plane Cremona group. In *Algebraic geometry in East Asia—Seoul 2008*, volume 60 of *Adv. Stud. Pure Math.*, pages 1–49. Math. Soc. Japan, Tokyo, 2010.

[4] W. Duke. The dimension of the space of cusp forms of weight one. *Internat. Math. Res. Notices*, (2):99–109, 1995.

[5] A. Fröhlich, editor. *Algebraic number fields: L-functions and Galois properties*. Academic Press [Harcourt Brace Jovanovich Publishers], London, 1977.

[6] Albrecht Fröhlich. *Central extensions, Galois groups, and ideal class groups of number fields*, volume 24 of *Contemporary Mathematics*. American Mathematical Society, Providence, RI, 1983.

[7] Daniel Gorenstein. *Finite groups*. Chelsea Publishing Co., New York, second edition, 1980.

[8] Jing Long Hoelscher. Ray class groups of quadratic and cyclotomic fields. *Int. J. Number Theory*, 6(5):1169–1182, 2010.

[9] I. Martin Isaacs. *Character theory of finite groups*. AMS Chelsea Publishing, Providence, RI, 2006. Corrected reprint of the 1976 original [Academic Press, New York; MR0460423].

[10] Stéphane R. Louboutin. Upper bounds for residues of Dedekind zeta functions and class numbers of cubic and quartic number fields. *Math. Comp.*, 80(275):1813–1822, 2011.

[11] Hyunsuk Moon. The number of monomial mod p Galois representations with bounded conductor. *Tohoku Math. J. (2)*, 55(1):89–98, 2003.

[12] L. B. Pierce. The 3-part of class numbers of quadratic fields. *J. London Math. Soc. (2)*, 71(3):579–598, 2005.

[13] Jean-Pierre Serre. *Local fields*, volume 67 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1979. Translated from the French by Marvin Jay Greenberg.

[14] Siman Wong. Automorphic forms on GL(2) and the rank of class groups. *J. Reine Angew. Math.*, 515:125–153, 1999.